

Instructions: Show all work. Answers without work required to obtain the solution will not receive full credit. Some questions may contain multiple parts: be sure to answer all of them. Give exact answers unless specifically asked to estimate.

- Solve the second-order ODEs for the general solution.
 - $2y'' - y' - y = 0$
 - $y'' - 2y' + 2y = 0$
 - $y'' - 18y' + 81y = 0$
- The table below gives the solution to the second order constant coefficient homogeneous equation, and the forcing function $F(x)$ or $F(t)$. Determine the Ansatz (particular solution, y_p) for the method of undetermined coefficients in each case.

	y_1	y_2	$F(x)$ or $F(t)$	Ansatz
a.	e^{-2x}	e^{3x}	$2 \sin 3x$	
b.	$e^{-x} \cos x$	$e^{-x} \sin x$	$e^x \sin x$	
c.	$e^{-x/2} \sin\left(\frac{\sqrt{3}}{2}x\right)$	$e^{-x/2} \cos\left(\frac{\sqrt{3}}{2}x\right)$	$e^x + 7$	
d.	e^{-t}	1	$t + e^{-t}$	
e.	$\sin t$	$\cos t$	$\cos^2 t$	

- What is the difference between the natural frequency of the system, and a quasi-frequency? How is each obtained?
- What conditions are needed in a forced oscillation system to achieve beats?
- Use the method of reduction of order to solve $(1 - x^2)y'' - 2xy' + 2y = 0$, given $y_1(x) = x$.
- Set up the differential equation to solve the spring-mass problem with a 12 lbs. weight that stretches a spring 6 in. and a dashpot that provides 3 lbs. of resistance for every ft/s of velocity. The weight is pulled from an additional one foot from equilibrium and then released from rest.
 - Is the system undamped, underdamped, critically damped or overdamped?
 - Solve for an equation for the position of the mass at any time t .
 - State the period (or quasi-period), amplitude and phase shift.
 - What is the behavior of the system as $t \rightarrow \infty$?
- Use the method of variation of parameters to find the particular solution to $y'' + 6y' + 9y = 4e^{2t} + e^{-t}$.


8. Use the method of undetermined coefficients to find the particular solution to $y'' + 6y' + 9y = 4e^{2t} + e^{-t}$.

9. Use the method of undetermined coefficients to find the particular solution to $2y'' + 3y' + y = t^2 + 3 \sin t, y(0) = 0, y'(0) = 1$.


10. Use the method of variation of parameters to find the particular solution to $y'' - 2y' + y = \frac{e^t}{1+t^2}$.

11. Below are the graphs of solutions to forced spring problems. Determine if the solution models resonance or beats (or neither). Explain your reasoning.

a. $y(t) = \cos(t) - \sin(t) + t \sin t$



b. $y(t) = 3 \sin 6t + 2 \cos(7t)$



12. Sketch a graph of what an overdamped spring system looks like.

13. For each of the solutions below to a forced oscillation system, state i) the transient or steady state solution, ii) whether the system is undamped, underdamped, critically damped or overdamped, and iii) if resonance or beats occurs.

a. $y(t) = e^{-t}(c_1 \cos 5t + c_2 \sin 5t) + 5 \cos 4t + 4 \sin 4t$

b. $y(t) = c_1 e^{-t} + c_2 e^{-2t} + \sin 3t$

c. $y(t) = c_1 \cos 2t + c_2 \sin 2t + \frac{1}{6}t \cos 2t$

14. Use Abel's Theorem to find the value of the Wronskian for $y'' + 2xy' + 8y = 0$.
15. Find the Wronskian for $\{t^2, t^2 \ln t\}$.
16. A mass of 100 g stretches a spring 5 cm. If the mass is set in motion from its equilibrium position with a downward velocity of 10 cm/sec, and there is no damping.
- Determine the position y of the mass at any time t .
 - When does the mass first return to equilibrium? (i.e. when is $y=0$?)
 - State the period, amplitude and phase shift.
17. A 400 gal tank initially contains 100 gal of brine containing 50 lbs of salt. Brine containing 1 lbs of salt per gal enters the tank at a rate of 5 gal/s, and the well-mixed brine flows out of the tank at the rate of 3 gal/s.
- Find an equation for the amount of salt in the tank at time t .
 - How much salt will the tank contain when the tank is completely full?
 - What is the concentration in the salt at time t ?
 - What is the limiting concentration of salt in the tank as $t \rightarrow \infty$?
 - At what time t is the concentration equal to 90% of its limiting value?
18. Suppose that the temperature of a cup of coffee obeys Newton's law of cooling. If the coffee has a temperature of 200° when freshly poured, and 1 minute later cooled to 180° in a room at 72° , determine when the coffee reaches a temperature of 120° (in minutes).
19. Xanax takes about 11.2 hours after peak levels to reach 50% of that level in the blood stream. If a drug test can read levels at 5% of peak levels for a single dose, how long will it take to clear the system according to the test? Write a differential equation to solve the problem.