



3. Evaluate the flux integral  $\int_S \vec{F} \cdot \vec{N} dS$  where  $\vec{F} = 2z\hat{i} - 4x\hat{j} + \hat{k}$  for the surface  $S: z = 12 - 3x - 4y$  in the first octant. (12 points)

4. Use the divergence theorem to evaluate  $\int_S \vec{F} \cdot \vec{N} dS$  for  $\vec{F}(x, y, z) = (2xy - 1)\hat{i} + (3yz + 2)\hat{j} + xz\hat{k}$  for the closed surface bounded by the cylinder  $x^2 + y^2 = 4$ ,  $z = 4$ , and the coordinate planes. (14 points)

5. Evaluate  $\int_C \vec{F} \cdot d\vec{r}$  using Stokes' Theorem for  $\vec{F}(x, y, z) = z\hat{i} + x\hat{j} + xyz\hat{k}$  for  $S: z = x^2 - y^2$ ,  $0 \leq x \leq 1, 0 \leq y \leq 1$ . (12 points)

6. Write an integral for the arc length of the curve  $\vec{r}(t) = \sqrt[3]{t^2}\hat{i} + \sqrt{t^3}\hat{j} + t\hat{k}$  on the interval  $[1, 4]$ . Evaluate it numerically. (6 points)

7. Find the directional derivative of the function  $w = \arcsin xyz$  at the point  $(1, \frac{1}{2}, 1)$  in the direction of  $\vec{v} = \langle -1, 3, -2 \rangle$ . In what direction is the directional derivative a maximum? (10 points)

8. Use the Fundamental Theorem of Line Integrals to evaluate  $\int_C (2xy + 1)dx + (x^2 - y)dy$  on the line segment from  $(0,2)$  to  $(4, -2)$ . (10 points)

Part 2: In this section you will record your answers on paper along with your work. After scanning, submit them to a Canvas dropbox as directed. These questions will be graded by hand.

9. Find an equation of the tangent plane to the surface  $\vec{r}(u, v) = u \cos v \hat{i} + u \sin v \hat{j} + u\hat{k}$  at the point  $u = 3, v = \frac{\pi}{4}$ . (8 points)

10. Find the unit tangent vector for  $\vec{r}(t) = (\sqrt{t^3} - 4)\hat{i} + (t^2 + 1)\hat{j}$ . (8 points)

11. Find the equation of the tangent plane to the surface  $xyz + z^{2/3} = y^2$  at  $(0,1,1)$ . (8 points)

12. Use Green's Theorem to evaluate  $\int_C xydx + x^2dy$  on the path described by the boundary of the graphs  $y = x^2, y = \sqrt{x}$  oriented counterclockwise. (12 points)

13. Evaluate the surface integral  $\int_S f(x,y,z)dS$  for  $f(x,y,z) = x^2z^2$ ,  $S$ : on the cone  $z^2 = x^2 + y^2$ , between the planes  $z = 1, z = 3$ . [Hint: converting to cylindrical/polar will help.] (12 points)

## Cylindrical

$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta \\z &= z \\x^2 + y^2 &= r^2 \\\tan^{-1}\left(\frac{y}{x}\right) &= \theta\end{aligned}$$

## Spherical

$$\begin{aligned}x &= \rho \cos \theta \sin \phi \\y &= \rho \sin \theta \sin \phi \\z &= \rho \cos \phi \\x^2 + y^2 + z^2 &= \rho^2 \\\tan^{-1}\left(\frac{y}{x}\right) &= \theta \\\cos^{-1}\left(\frac{z}{\sqrt{x^2 + y^2 + z^2}}\right) &= \phi \\x^2 + y^2 &= \rho^2 \sin^2 \phi = r^2\end{aligned}$$

## Dels

$$\begin{aligned}\frac{\partial}{\partial x} &= \text{partial derivative with respect to } x \\\nabla f &= \text{grad } f \\\nabla^2 f &= \nabla \cdot (\nabla f) = \text{Laplacian of } f \\\nabla \cdot \vec{F} &= \text{div } \vec{F} \\\nabla \times \vec{F} &= \text{curl } \vec{F}\end{aligned}$$

## Misc

$$\begin{aligned}ds &= \|\vec{r}'(t)\| dt \\\kappa &= \frac{\|\vec{r}' \times \vec{r}''\|}{\|\vec{r}'\|^3} = \frac{1}{R} \\\vec{N} &= \nabla F = \vec{r}_u \times \vec{r}_v\end{aligned}$$